Family *Liliaceae Tulipa saxatilis* Sieber ex Spreng.

Geographical distribution: Crete, Karpathos, E.Aegean Islands & SW Turkey

Endemism:

Habitat: ledges & crevices of calcareous cliffs, rocky slopes & screes, stream banks in scrub, flat clayey areas, cultivated, fallow & abandoned fields (alt. 0-900 m)

Short description: Bulbous plant with simple bulbs $20-35 \times 15-30$ mm, covered with brown tunico, coriaceous, with a few straight hairs inside. Leaves 2-3(-4) up to 38 cm, oblong-lanceolate, glaucous-green. Flowers 1-2(-4), perianth pink to lilac-purple at tip, with a distinct white-edged, yellow basal blotch inside. Anthers 4,5-7 mm, filaments 8-17 mm, both golden-yellow.

Flowering period: March-May

Common names (in Crete, Greece): toulipa

Uses: ornamental plant

Conservation status (for threatened species):

Protection status (for threatened species): Greek Presidential Decree 67/1981

Code(s) of specimens in the Herbarium of MAICh: 3850

References:

Briggs M. 1990. First record of Tulipa goulimyi (Liliaceae) on Crete. Ann. Musei Goulandris 8:41-44.

Georgiou K & Delipetrou P. 2001. Threatened and Endemic plant species of South Greece. Region of Crete - Natural History Museum of Crete Grey Wilson C. 1998. A paradise for flowers. Garden (London) 123(3): 170-175 Protopapadakis, E.E. and Yannitsaros, A. 1994. Identification by isozymes of nine populations of Tulipa from Greece. Journal of Horticultural Science. 69(1) 11-14.

Sfikas, G. 1987. Wild Flowers of Crete. Efstathiadis & Sons S.A., Athens. 310 p..

Turland, N.J., Chilton L. & J.R. Press. 1993. Flora of the Cretan Area. Annotated Checklist & Atlas. The Natural History Museum. St Edmundsbury Press, Bury St. Edmunds, Suffolk, England.

Tutin, T. G. et al. (eds). Flora Europaea Vol. 5. p. 29. Cambridge University Press.

Iatrou G. 1986. Contribution to the study of the endemic flora of the Peloponnese. Ph.D. Thesis. University of Patras (in Greek).